

Hadronic final states in high- p_T QCD with the CMS detector

Suvadeep Bose


University of Nebraska Lincoln


(On behalf of CMS collaboration)


- ❑ Introduction
- ❑ Inclusive Jet production
 - at 7 TeV, **at 8 TeV**
 - AK5/AK7 ratio
- ❑ Dijet Production
 - Differential cross section at 7 TeV, **at 8 TeV**
 - Dijet mass and jet substructure
- ❑ Multi-jet Production
 - Colour coherence
 - 3-Jet Mass cross section
 - 3-jet to 2-jet cross section Ratio
 - Measurement of α_s


All data from published results are posted on the Durham database:
<http://hepdata.cedar.ac.uk/>


For all public results in CMS Standard Model Physics:
<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSMP>



















**Reality is
more complex**



- ❑ It is interesting
 - very rich theory: deserves exploration and understanding
- ❑ It is inevitable
 - hadron collisions: QCD is always present
- ❑ Important background for new physics searches
 - enormous cross section: QCD can hide many possible signals of new physics
- ❑ Introduces uncertainties on other measurements
 - e.g. uncertainties on the PDFs affect the Higgs properties
- ❑ With LHC data
 - probing new territory



experimental access to an enormous phase space → stringent test of QCD

CMS: the detector

Total weight
14000 t
Diameter 15 m
Length 28.7 m

ECAL 76k scintillating
PbWO₄ crystals

HCAL Scintillator/brass
Interleaved ~7k ch




**MUON
ENDCAPS**
473 Cathode Strip Chambers (CSC)
432 Resistive Plate Chambers (RPC)

3.8T Solenoid


IRON YOKE



Thanks LHC for Fantastic 3 years!



Particle-Flow (PF) algorithms use:
e, μ , γ , charged & neutral hadrons
as building blocks for
jets, b-jets, τ 's, ν (miss E_T)




Fixed cone algorithms:


- ✧ Iterative Cone (CMS) / JetClu (ATLAS)
- ✧ Seedless Infrared Safe Cone (SISCone)

Successive recombination algorithms:

$$d_{ij} = p_{T,i}^{2p} \quad d_{ij} = \min(p_{T,i}^{2p}, p_{T,j}^{2p}) \frac{\Delta R_{ij}^2}{D^2}$$



kT jet Cone jet





p=1	->	k _T jet algorithm
p=0	->	CA jet algorithm
p=-1	->	"Anti-k _T " jet algorithm


- Soft particles will first cluster with hard particles before among themselves
- Almost a cone jet near hard partons
- No merge/split

- Jet Energy Correction is necessary to measure the correct energy spectrum of Jets
- The main three type of corrections required are **Offset** (pile up subtraction), **Relative** (for η dependent response) and **Absolute** (for p_T dependent response) + Residual corrections. [JINST 6 (2011) P11002]
 - Offset \rightarrow **Subtracting**
 - Relative \rightarrow **Dijet balance**
 - Absolute \rightarrow **γ + jet and Z + jet (p_T balance, MPF)**

JEC (Jet Energy Correction) uncertainty $\sim 1\%$ for central high- p_T jets




CMS DP-2013/011




Inclusive Jet Production

- Inclusive Jet Cross Section @ 7 TeV
[QCD-11-004]
[Phys. Rev. D 87 \(2013\) 112002](#)
- Inclusive Jet Cross Section @ 8 TeV
[SMP-12-012, FSQ-12-031]
- Inclusive jet AK5/AK7 cross section ratio @ 7 TeV [SMP-13-002]




UA2: PLB 118 (1982)
CERN pp-bar $\sqrt{s} = 540$ GeV





.. after 30 years

Fig. 6. Inclusive jet production cross section. The solid line (ref. [6]) uses $\Lambda = 0.5$ GeV while $\Lambda = 0.15$ GeV would bring the calculated rates in better agreement with the data. However various uncertainties preclude a determination of Λ from the data [13].




- Inclusive jet production probes the dynamics of QCD
- counting the number of jets as a function of rapidity and p_T stringent test of QCD
- PDFs, strong coupling constant, perturbative calculations



- Double differential Inclusive jet cross-section measured from p_T 0.1 to 2 TeV
- Measured cross sections agree with the predictions of perturbative QCD at next-to-leading order obtained with five different PDF sets
- Theoretical and experimental uncertainties are comparable, even at the limits of the experimental phase space

SMP-12-012
FSQ-12-031



❑ Measurement at 8 TeV up to $|y| = 4.7$

❑ **O(14)** magnitude in cross section

❑ Comparisons to pQCD NLO \otimes NP

[NP: corrections for non-perturbative effects (MPI and hadronization):

20% (~ 100 GeV) \rightarrow 1% (~ 2.5 TeV)]

SMP-12-012





Experimental uncertainty:



- JES (12%-30%),
- Luminosity (4.4%)
- Unfolding (1%-10%)
- Total: 15%-40%

Theory uncertainty:

- PDF (5%-30%)
- Scale (5%-40%)



PDF uncertainty for CT10 in outer bins **100%**





- ❑ Data over theory compared to ratio with other PDF sets for **CT10**
- ❑ The theory predictions are computed for five different PDF sets, viz. **ABM11**, **HERA1.5**, **CT10**, **MSTW2008**, **NNPDF2.1**
- ❑ In the central rapidity region ($0.0 < |y| < 0.5$) different theory predictions are in agreement with data except **ABM11**
- ❑ The fluctuations are covered by total theoretical and experimental uncertainty bands

- ❑ Motivation: AK5/Ak5 ratio gives insight to QCD effect beyond fixed order
- ❑ Measurement at $\sqrt{s}=7$ TeV with different jet sizes $R=0.5$ (AK5), 0.7 (AK7)
- ❑ Ratio of cross sections $R(0.5, 0.7)$ vs p_T and rapidity




SMP-13-002




- ❑ Several systematic uncertainties cancel in ratio
- ❑ The ratio gradually increases towards unity with increasing Jet- p_T .
- ❑ Powheg(NLO+PS) prediction has the describes the data best

Dijet Production

- Dijet Differential Cross Section
@ 7 TeV [QCD-11-004]
[Phys. Rev. D 87 \(2013\) 112002](#)
- Dijets and V+jets, jet mass and
substructure at 7 TeV [SMP-12-019]
[JHEP 05 \(2013\) 090](#)





QCD-11-004



- Reach up to $M_{jj} \sim 5.5 \text{ TeV}$
- Complementary to Inclusive jets
- Agreement with pQCD@NLO \otimes NP

- ❑ Differential distributions in jet mass for inclusive dijet events, defined through the anti- k_T algorithm for a size parameter of 0.7 for jets groomed through **filtering**, **trimming**, and **pruning**.

SMP-12-019




The intrinsic stability of these algorithms to pileup effects contribute to a more rapid and widespread use of these techniques in future high-luminosity runs at the LHC.

- Better agreement at larger jet masses.
- Trimming and pruning algorithms provide an important benchmark for their use in searches for massive particles.

➤ More details in [Ivan Marchesini's](#) talk on Jet substructure (Today at 3 pm)

MultiJet Production


- Colour Coherence [SMP-12-010]
- 3-Jet Mass cross section
[SMP-12-027]
- 3/2 Inclusive Jet Cross section Ratio
[QCD-11-003] [arXiv:1304.7498](https://arxiv.org/abs/1304.7498)
- Measurement of α_s
[SMP-12-027, QCD-11-003]




- ❑ In QCD color coherence effects are due to the interference of soft gluon radiation emitted along color connected partons
- ❑ In LO model with FSR radiation the 3rd jet tends to be between second jet and proton remnant
- ❑ Measure the angular distribution of softer 3rd jet around the 2nd highest- p_T jet in the event

$$\beta = \tan^{-1} \left[\frac{\text{sign}(\eta_2) \Delta\phi_{32}}{\Delta\eta_{32}} \right]$$

- $\beta=0$: 3rd jet between 2nd & closest proton remnant
 $\beta=\pi$: 3rd jet between 2nd & far most proton remnant





- ❑ Ideally, $e^+ e^-$ collider is the best place to do the measurement
 - No color interference from the initial state
- ❑ In pp , both initial and final states have color constituents
 - Complicate the signatures
 - Comparison with MC is crucial
- ❑ Compare data to event generators with different color coherence implementations



- Data are unfolded to the particle level
- The data exhibit a clear enhancement of events compared with the PYTHIA and MADGRAPH generators near the event plane ($\beta = 0$) and a suppression in the transverse plane ($\beta = \pi/2$)

- Comparisons of the β distributions to various MC predictions
 - Herwig++ describes the data β distributions best

- Data clearly support larger color coherence effects in Pythia 6




SMP-12-010



Measurement of double differential cross section: $d^2\sigma/dm_3 dy_{\max}$

- sensitivity to PDFs and α_s
- $m_3^2 = (p_1 + p_2 + p_3)^2$ $|y_{\max}| = \max(|y_1|, |y_2|, |y_3|)$ $Q = m_3/2$



- Require jet $p_T > 100$ GeV
- Regions: $|y_{\max}| < 1$ and $1 < |y_{\max}| < 2$ - reach up to $m_3 \sim 3$ TeV
- Agreement with pQCD @ NLOxNP (*NP correction 8% -> 1%*)
 - Deviations observed with NLO + ABM11 PDFs



SMP-12-027




QCD-11-003



- ❑ Cross section ratio R_{32} :
 - inclusive 3-jet over 2-jet production
 - sensitive to α_s
- ❑ Multiple alternative phase-space options
 - depending on the cut imposed on the 3rd jet p_T
 - expressed vs. different observables
 - **measuring the α_s** : vital to reduce scale uncertainty

$$R_{32} = \frac{\sigma_3}{\sigma_2} = \frac{\sigma(pp \rightarrow n \text{ jets} + X; n \geq 3)}{\sigma(pp \rightarrow n \text{ jets} + X; n \geq 2)} \quad \text{vs} \quad \langle p_{T1,2} \rangle = \frac{p_{T1} + p_{T2}}{2}$$



SMP-12-027
QCD-11-003
[arXiv:1304.7498](https://arxiv.org/abs/1304.7498)

- Extract α_s from the R_{32} and 3-jet mass cross section measurements
 - Results are comparable with world average $\alpha_s(M_Z) = 0.1184 \pm 0.0007$
 - For the first time probing the $> 1 \text{ TeV}$ scale, reaching up to $\sim 1.5 \text{ TeV}$
 - Dominated by theoretical uncertainties (PDF and scale)
- R_{32} : $\alpha_s(M_Z) = 0.1148 \pm 0.0014 \text{ (exp.)} \pm 0.0018 \text{ (PDF)}^{+0.0050}_{-0.0000} \text{ (scale)}$
- 3-jet mass: $\alpha_s(M_Z) = 0.1160^{+0.0025}_{-0.0023} \text{ (exp, PDF, NP)}^{+0.0068}_{-0.0021} \text{ (scale)}$

- ❑ Significant ongoing effort to improve our understanding of QCD
 - both experimental and theoretical
 - rich QCD programs pursued LHC
- ❑ Large datasets available
 - LHC has provided access to a huge phase space
 - will take a long time to analyze and digest all the data on tape
- ❑ Much recent progress
 - jet data have considerable impact on gluon and u/d quark PDFs
 - measurements of α_s at the TeV scale for the first time
- ❑ Comments on the theoretical tools
 - in many areas the exp. precision reached makes the NLO predictions insufficient: NNLO needed for further progress!!
 - with some tuning of the parameters, the LO ME or NLO interfaced with PS models provide good description of the data